

# इंटरनेट

# मानक

## Disclosure to Promote the Right To Information

Whereas the Parliament of India has set out to provide a practical regime of right to information for citizens to secure access to information under the control of public authorities, in order to promote transparency and accountability in the working of every public authority, and whereas the attached publication of the Bureau of Indian Standards is of particular interest to the public, particularly disadvantaged communities and those engaged in the pursuit of education and knowledge, the attached public safety standard is made available to promote the timely dissemination of this information in an accurate manner to the public.

“जानने का अधिकार, जीने का अधिकार”

Mazdoor Kisan Shakti Sangathan

“The Right to Information, The Right to Live”

“पुराने को छोड़ नये के तरफ”

Jawaharlal Nehru

“Step Out From the Old to the New”

IS 7203 (1973): Casein Hydrolysate (Acid Digested),  
Microbiological Grade [FAD 15: Food Hygiene, Safety  
Management and Other Systems]



“ज्ञान से एक नये भारत का निर्माण”

Satyanarayan Gangaram Pitroda

“Invent a New India Using Knowledge”



“ज्ञान एक ऐसा खजाना है जो कभी चुराया नहीं जा सकता है”

Bhartrhari—Nitiśatakam

“Knowledge is such a treasure which cannot be stolen”



BLANK PAGE



*Indian Standard*  
SPECIFICATION FOR  
CASEIN HYDROLYSATE ( ACID DIGESTED ),  
MICROBIOLOGICAL GRADE

UDC 577.152:663.1



© Copyright 1974

INDIAN STANDARDS INSTITUTION  
MANAK BHAVAN, 9 BAHADUR SHAH ZAFAR MARG  
NEW DELHI 110001

Price Rs. 4.00

September 1974

Revised Price

Gr 3

# Indian Standard

## SPECIFICATION FOR CASEIN HYDROLYSATE ( ACID DIGESTED ), MICROBIOLOGICAL GRADE

Food Hygiene, Sampling and Analysis Sectional Committee, AFDC 36

### Chairman

MAJ-GEN M. S. BOPARAI

### Representing

Directorate General, Armed Forces Medical  
Services ( Ministry of Defence ), New Delhi

### Members

AGRICULTURAL MARKETING AD- VISER TO THE GOVERNMENT OF INDIA	Directorate of Marketing & Inspection ( Ministry of Agriculture ), Faridabad
SHRI T. V. MATHEW ( <i>Alternate</i> )	
SHRI V. N. AMBLE	Institute of Agricultural Research Statistics ( ICAR ), New Delhi
SHRI K. S. KRISHNAN ( <i>Alternate</i> )	
SHRI K. BALASUBRAMANIAN	Public Analyst, Government of Tamil Nadu, Madras
LT S. L. CHADHA	Health Department, Municipal Corporation of Delhi
DR A. G. AJWANI ( <i>Alternate</i> )	
DR G. C. DAS	Corporation of Calcutta
DR P. K. DATTA	All India Institute of Hygiene and Public Health, Calcutta
SHRI SUKUMAR DE	National Dairy Research Institute ( ICAR ), Karnal
DR C. A. MULAY ( <i>Alternate</i> )	
DIRECTOR	Central Food Laboratory, Calcutta
EXECUTIVE HEALTH OFFICER	Bombay Municipal Corporation
MUNICIPAL ANALYST ( <i>Alternate</i> )	
HEALTH OFFICER	Corporation of Madras
DR ( SMT ) S. KHOSLA	Department of Health & Family Planning, Government of Punjab, Chandigarh
DR P. K. KYMAL	Food & Nutrition Board ( Ministry of Agriculture ), New Delhi
DR R. C. SINHA ( <i>Alternate</i> )	
SHRI Y. N. LAKSHMAN	Defence Food Research Laboratory ( Ministry of Defence ), Mysore
DR G. N. VERMA ( <i>Alternate</i> )	
SHRI F. G. T. MENEZES	Directorate of Sugar & Vanaspati ( Ministry of Agriculture ), New Delhi
SHRI D. VENKATAPPAIA ( <i>Alternate</i> )	

( Continued on page 2 )

© Copyright 1974

INDIAN STANDARDS INSTITUTION

This publication is protected under the *Indian Copyright Act* ( XIV of 1957 ) and reproduction in whole or in part by any means except with written permission of the publisher shall be deemed to be an infringement of copyright under the said Act.

(Continued from page 1)

<i>Members</i>	<i>Representing</i>
PUBLIC ANALYST (FOOD AND WATER)	Government of West Bengal, Calcutta
DR T. N. RAMACHANDRA RAO	Central Food Technological Research Institute (CSIR), Mysore
SHRI C. T. DWARKANATH (Alternate)	
DR RANJIT SEN	Directorate General of Health Services (Ministry of Health & Family Planning), New Delhi
SENIOR COMMERCIAL OFFICER (CATERING)	Chief Catering Officer, Northern Railway, New Delhi
COL K. N. SHARMA	Quarter Master General's Branch, Army Headquarters
LT-COL O. P. KAPUR (Alternate)	
DR S. B. SINGH	Public Analyst, Government of Uttar Pradesh, Lucknow
DR M. C. SWAMINATHAN	Central Committee for Food Standards, New Delhi
SHRI D. S. CHADHA (Alternate)	
SHRI P. C. VIN	The Coca-Cola Export Corporation, New Delhi
SHRI J. D. CONTRACTOR (Alternate)	
DR HARI BHAGWAN, Director (Agri & Food)	Director General, ISI (Ex-officio Member)

*Secretary*

SHRI SOHRAB  
Assistant Director (Agri & Food), ISI

Media for Microbiological Tests Subcommittee, AFDC 36 : 7

*Convener*

DR RANJIT SEN	Serologist to the Government of India (DGHS), Calcutta
---------------	--

*Members*

DR A. N. BOSE DIRECTOR	Bengal Immunity Co Ltd, Calcutta King Institute, Madras
DR D. V. MURTHY (Alternate)	
DR A. K. GHOSH	Cholera Research Centre (ICMR), Calcutta
HEAD, DIVISION OF BIOLOGICAL PRODUCTS	Indian Veterinary Research Institute (ICAR), Izatnagar
DR A. P. JOSHI	Vallabhai Patel Chest Institute, Delhi
DR (SMT) V. BAJAJ (Alternate)	
DR A. D. MUKHERJEE	Bengal Chemical and Pharmaceutical Works Ltd, Calcutta
DR A. N. RAI CHOWDHURI	Central Research Institute, Kasauli
DR T. N. RAMACHANDRA RAO	Central Food Technological Research Institute (CSIR), Mysore
DR B. RANGANATHAN	National Dairy Research Institute (ICAR), Karnal
DR N. S. SUBBA RAO	Indian Agricultural Research Institute (ICAR), New Delhi
DR M. V. SANT	Haffkine Institute, Bombay

# *Indian Standard*

## SPECIFICATION FOR CASEIN HYDROLYSATE ( ACID DIGESTED ), MICROBIOLOGICAL GRADE

### 0. F O R E W O R D

**0.1** This Indian Standard was adopted by the Indian Standards Institution on 26 November 1973, after the draft finalized by the Food Hygiene, Sampling and Analysis Sectional Committee had been approved by the Agricultural and Food Products Division Council.

**0.2** Unless the ingredients used in media for microbiological analysis are of uniform quality, the results obtained might be erroneous and might be unreliable. Since the media used in different laboratories often differ greatly in their quality, the results of microbiological analysis at different laboratories cannot be compared. Therefore, with a view to unifying the practices of different laboratories dealing with microbiological work and providing guidance to the indigenous manufacturers regarding the quality, it was decided to bring out a series of Indian Standard specifications for ingredients commonly used in media for microbiological analysis.

**0.3** Casein hydrolysate ( acid digested ) microbiological grade, is used for microbiological work especially for growing whooping-cough organism in large quantities for production of vaccine. It is a yellowish white fine powder, highly soluble in water but insoluble in alcohol, ether and chloroform.

**0.4** For the purpose of deciding whether a particular requirement of this standard is complied with, the final value, observed or calculated, expressing the result of a test, shall be rounded off in accordance with IS:2-1960\*. The number of significant places retained in the rounded off value should be the same as that of the specified value in this standard.

---

### 1. SCOPE

**1.1** This standard prescribes the requirements and the methods of tests for casein hydrolysate ( acid digested ) microbiological grade.

---

\*Rules for rounding off numerical values ( revised ).

## 2. REQUIREMENTS

2.1 A one percent solution in water shall show colourless to light straw yellow liquid having a pH between 6.0 and 7.0.

2.2 It shall be able to support growth of *Bordetella pertussis* when tested according to method shown in Appendix A.

2.3 Iron — It shall pass the test prescribed in Appendix B.

2.4 The material shall also conform to the requirements given in Table 1.

TABLE 1 REQUIREMENTS FOR CASEIN HYDROLYSATE

SL No.	CHARACTERISTIC	REQUIREMENT	METHOD OF TEST		
			Ref to Appendix of this Standard	Ref to Clauses of IS: 6854-1972*	Ref to Appendix of IS: 7128-1973†
(1)	(2)	(3)	(4)	(5)	(6)
i)	Moisture content, percent by mass, <i>Max</i>	7	—	4	—
ii)	Total nitrogen, percent by mass, <i>Min</i>	8	—	9	—
iii)	Amino nitrogen, percent by mass, <i>Min</i>	6	—	8	—
iv)	Sodium chloride, percent by mass, <i>Min</i>	30	—	11	—
v)	Ash, percent by mass, <i>Min</i>	35	—	6	—
vi)	Ether soluble extract, percent by mass, <i>Max</i>	0.1	C	—	—
vii)	Phosphorus (as P), percent by mass, <i>Max</i>	0.9	—	—	E

\*Method of sampling and test for ingredients used in media for microbiological work.

†Specification for proteose peptone, microbiological grade.



### 3. PACKING, STORAGE AND MARKING

**3.1 Packing** — The material shall be securely packed in well-filled amber coloured moisture proof wide mouth containers with tight fitting lids.

**3.2 Storage** — The material shall be stored in a cool and dry place.

**3.3 Marking** — Each container shall be marked legibly to give the following information:

- a) Name of the material including the words 'Microbiological grade',
- b) Name and address of the manufacturer,
- c) Minimum net content,
- d) Batch or code number, and
- e) The label should indicate quantity of sodium chloride, total nitrogen and phosphate. It shall also indicate that the material passes the prescribed test for iron.

**3.3.1** The container may also be marked with the ISI Certification Mark.

**NOTE** — The use of the ISI Certification Mark is governed by the provisions of the Indian Standards Institution (Certification Marks) Act and the Rules and Regulations made thereunder. The ISI Mark on products covered by an Indian Standard conveys the assurance that they have been produced to comply with the requirements of that standard under a well-defined system of inspection, testing and quality control which is devised and supervised by ISI and operated by the producer. ISI marked products are also continuously checked by ISI for conformity to that standard as a further safeguard. Details of conditions under which a licence for the use of the ISI Certification Mark may be granted to manufacturers or processors, may be obtained from the Indian Standards Institution.

### 4. SAMPLING

**4.1** The representative samples of the material shall be drawn according to the method prescribed in 3 of IS:6854-1972\*.

### 5. TESTS

**5.1** Tests shall be carried out by the methods prescribed in 2.2 and in col 4, 5 and 6 of the Table 1.

**5.2 Quality of Reagents** — Unless specified otherwise, pure chemicals and distilled water ( *see* IS:1070-1960† ) shall be employed in the tests.

**NOTE** — 'Pure chemicals' shall mean chemicals that do not contain impurities which affect the experimental results.

\*Methods of sampling and test for ingredients used in media for microbiological work.

†Specification for water, distilled quality ( *revised* ).

## APPENDIX A

( Clause 2.2 )

## TEST FOR ABILITY TO SUPPORT GROWTH

## A-1. GROWTH SUPPORTING PROPERTIES

**A-1.1 Strain** — *Bordetella pertussis* strain No. 134 or 509 ( Dutch strain ) or any other suitable strain in phase I approved by an International authority. Maintain the strain in such a way that the organism is retained in phase I.

## A-2. BORDET-GENGOU MEDIUM

**A-2.1 Base Medium** — Prepare the medium with 12.5 percent peeled and cleaned potato slices, 0.45 percent sodium chloride, 1.0 percent glycerol, 1.0 percent proteose-peptone ( see IS : 7128-1973\* ) and 2.25 percent agar ( see IS : 6850-1973† ) in distilled water. Cut into thin slices the peeled and cleaned potatoes and boil with sodium chloride in one-fourth the amount of distilled water until the potato slices fall to pieces. Make up the water lost during boiling and filter through linen and adjust the reaction to pH 7.0. Dissolve the agar separately by boiling in the remaining amount of distilled water and to this add the boiled and filtered potato extract, glycerol and proteose-peptone and mix well. Distribute into bottles and sterilize by steaming at 100°C for one hour followed by autoclaving at 115°C for 10 minutes.

**A-2.2 Complete Medium** — Prepare the complete medium with the 2 volumes of base medium ( A-2.1 ) and 1 volume of sterile defibrinated horse blood. Melt the base medium ( A-2.1 ) by steaming for one hour at 100°C mixing well during heating. Place in a water bath at 55°C until the temperature of the liquid base medium drops to about 65°C. Warm sterile defibrinated horse blood in the water bath at 55°C for 2 to 3 minutes and add to the liquid base medium. Mix well and pour at least 30 ml into sterile petri dishes of 10 cm diameter.

## A-3. TEST PROCEDURE

**A-3.1** Prepare medium with the following composition:

Casein hydrolysate ( Acid digested ), percent	1.4
Potassium chloride ( KCl ), percent	0.02

---

\*Specification for proteose-peptone, microbiological grade.

†Specification for agar, microbiological grade.

Monobasic anhydrous potassium phosphate ( $\text{KH}_2\text{PO}_4$ ), percent	0.05
Magnesium chloride ( $\text{MgCl}_2, 6\text{H}_2\text{O}$ ), percent	0.01
Nicotinic acid, percent	0.002
Glutathione, percent	0.001
Starch, percent	0.1
Distilled water	

Distribute in 100 ml amounts into Roux bottles and sterilize by autoclaving at  $115^\circ\text{C}$  for 20 minutes.

**A-3.2** Grow the strain of *B. pertussis* on a Bordet-Gengou plate ( **A-2.2** ) at  $35^\circ\text{C}$  for 72 hours and subculture on to another Bordet-Gengou plate. Incubate at  $35^\circ\text{C}$  for 48 hours and emulsify the growth in 5 ml of sterilized solution of 1 percent casein hydrolysate ( acid digested ). Inoculate 0.1 ml of the suspension into 100 ml of the medium as prepared in **A-3.1**. Incubate at  $35^\circ\text{C}$  for 40 to 48 hours taking care to lay the Roux bottle on its side so as to expose a large surface. Make two further serial subcultures into the medium as prepared in **A-3.1** using 1 ml as the inoculum for these serial subcultures. Each time incubate at  $35^\circ\text{C}$  for 40 to 48 hours exposing a large surface of the medium. Transfer the entire contents of the last subcultured and 40 to 48 hours incubated Roux bottle of medium ( **A-3.1** ) to suitable containers. Centrifuge at 2 500 rpm for 40 minutes. Wash the deposits with saline and pool into one container and centrifuge again. Suspend the deposit in 15 ml of 0.85 percent saline.

#### A-4. REQUIREMENTS

**A-4.1** The saline suspended material as obtained from **A-3.2** shall meet the following requirements:

- a) *Bacterial Density* — It shall have a bacterial density of more than  $2 \times 10^{10}$  organisms per ml when determined by the following procedure:

‘Not less than  $\frac{1}{2}$  dilution of the suspended material diluted with physiological saline should match, when inspected visually, with the WHO International Opacity Reference Preparation containing 10 units of opacity which is equivalent to  $1 \times 10^{10}$  organisms per ml.

- b) *Phase Variations* — It shall show profuse growth of *B. pertussis* when inoculated on Bordet-Gengou plate and no growth on ordinary nutrient agar medium.

## APPENDIX B

( Clause 2.3 )

### LIMIT TEST FOR IRON

#### B-1. REAGENTS

**B-1.1 Casein Hydrolysate Solution** — Dissolve 2.3 g casein hydrolysate in 50 ml of distilled water. Make up the volume to 100 ml in a volumetric flask.

**B-1.2 Hydrochloric Acid, Dilute** — Dilute 250 ml concentrated hydrochloric acid ( sp gr 1.18 ), reagent grade, to 1000 ml with distilled water.

**B-1.3 Standard Iron Solution** — Dissolve 0.173 g ferric ammonium sulphate, reagent grade, in 100 ml distilled water. Add 5 ml dilute hydrochloric acid and make up the volume to 1000 ml in a volumetric flask. This solution contains 0.02 mg iron in 1 ml.

**B-1.4 Ammonia Solution, Dilute** — Dilute 375 ml of ammonia solution ( sp gr 0.88 ), reagent grade to 100 ml with distilled water.

**B-1.5 Citric Acid** — 20 percent solution.

**B-1.6 Thioglycollic Acid**

#### B-2. PROCEDURE

**B-2.1** To three, 50 ml capacity Nessler cylinders, labelled Blank, Standard and Casein hydrolysate respectively, add 40 ml of distilled water to the blank, 2 ml of standard iron solution diluted to 40 ml with distilled water to the standard and 35 ml of casein hydrolysate solution to the casein hydrolysate cylinders.

**B-2.2** Add in succession to each cylinder, 2 ml of citric acid solution and 2 drops of thioglycollic acid, and mix well.

**B-2.3** Make the solutions alkaline to about pH 8 using pH test paper with dilute ammonia solution, make up the volumes to 50 ml with distilled water and mix well.

**B-2.4** Allow the solutions to stand for 5 minutes. The colour produced by casein hydrolysate should not be more intense than the colour produced by the standard iron solution. The blank should be free of any tinge of pink colour.

## APPENDIX C

[ *Table 1, Item (vi)* ]

### DETERMINATION OF ETHER SOLUBLE EXTRACT

#### C-1. REAGENT

C-1.1 Diethyl Ether — anhydrous.

#### C-2. PROCEDURE

C-2.1 Extract 2 g of the ground material in a continuous extraction apparatus with diethyl ether for 18 hours. Remove the ether by distillation, followed by heating the flask on a boiling water-bath. Dry the residue in an oven at  $110\text{ C} \pm 1^{\circ}\text{C}$  till the loss in mass between two successive weighings is less than 2 mg. Shake the residue with 2 to 3 ml of diethyl ether at room temperature, allow the residue to settle and decant the ether. Repeat the extraction until no more of the residue dissolves. Dry the flask again until the loss in mass between two successive weighings is less than 2 mg. Note the lowest mass.

#### C-3. CALCULATION

$$\text{Non-volatile ether extract, percent by mass} = \frac{100 (W_1 - W_2)}{W}$$

where

$W_1$  = mass in g of the flask with the non-volatile ether extract,

$W_2$  = mass in g of the flask with the ether insoluble residue after decantation, and

$W$  = mass in g of the material taken for the test.

**INDIAN STANDARDS**  
**ON**  
**FOOD HYGIENE, SAMPLING AND ANALYSIS**

IS:

- 2491-1972 Code for hygienic conditions for food processing units ( *first revision* )
- 5059-1969 Code for hygienic conditions for large scale biscuit manufacturing units and bakery units
- 5398-1969 Methods for estimation of thiamine (vitamin B<sub>1</sub>) in foodstuffs
- 5399-1969 Methods for estimation of riboflavin (vitamin B<sub>2</sub>) in foodstuffs
- 5400-1969 Methods for estimation of nicotinic acid (niacin) foodstuffs
- 5401-1969 Methods for detection and estimation of coliform bacteria in foodstuffs
- 5402-1969 Method for plate count of bacteria in foodstuffs
- 5403-1969 Method for yeast and mould count of foodstuffs
- 5404-1969 Code of practice for handling of food samples for microbiological analysis
- 5835-1970 Method for estimation of vitamin D in foodstuffs
- 5837-1970 Code for hygienic conditions for soft drinks manufacturing units
- 5838-1970 Method for estimation of vitamin C in foodstuffs
- 5839-1970 Code for hygienic conditions for manufacture, storage and sale of ice-creams
- 5886-1970 Method for estimation of carotenes and vitamin A ( retinol ) in foodstuffs
- 5887-1970 Methods for detection of bacteria responsible for food poisoning and food-borne diseases
- 6540-1972 Code for hygienic conditions for manufacturer and handling of ice for human consumption
- 6541-1972 Code for hygienic conditions for establishment and maintenance of midday school meal programmes
- 6542-1972 Code for hygienic conditions for fruit and vegetable canning units
- 6850-1973 Agar, microbiological grade
- 6851-1973 Meat extract, microbiological grade
- 6852-1973 Bile salts, microbiological grade
- 6853-1973 Peptone, microbiological grade
- 6854-1973 Methods of sampling and test for ingredients used in media for microbiological work
- 6968-1973 Code for hygienic conditions for *PAN* ( betel ) stalls and vendors
- 6969-1973 Code for hygienic conditions for handling and sale of refrigerated drinking water
- 7003-1973 Code for hygienic conditions for sago ( *SABOODANA* ) manufacturing units
- 7004-1973 Yeast extract, microbiological grade
- 7005-1973 Code for hygienic conditions for production, processing, transportation and distribution of milk

# PUBLICATIONS OF INDIAN STANDARDS INSTITUTION

## INDIAN STANDARDS

Over 7 500 Indian Standards covering various subjects have been issued so far. Of these, the standards belonging to the Agricultural and Food Products Group fall under the following categories:

Abattoir	Food additives
Alcoholic drinks	Food grain handling and storage
Animal feeds	Fruits and vegetables
Animal housing and equipment	Honey and by-products
Bakery and confectionery	Infant foods
Bee-keeping equipment	Laboratory animals
Beverages	Meat and meat products
Cereals, pulses and their products	Pest control equipment
Cocoa products	Pesticidal formulations
Coffee and its products	Pesticides, technical grade and general
Dairy equipment	Propagation materials
Dairy industry, layout plans	Regulated market yards
Dairy industry, methods of test	Sensory evaluation
Dairy laboratory apparatus	Spices and condiments
Dairy products	Starch derived products
Edible starch and starchy products	Sugars and by-products
Farm implements and machinery	Tea
Fish and fishery products	Tobacco products
Fish industry, sanitary conditions	Transport of live animals

## OTHER PUBLICATIONS

ISI Bulletin (Published Every Month)					
Single Copy	...	...	...	...	Rs 3-00
Annual Subscription	...	...	...	...	Rs 25-00
Standards: Monthly Additions					
Single Copy	...	...	...	...	Rs 0-30
Annual Subscription	...	...	...	...	Rs 3-00
Annual Reports (from 1948-49 Onwards)	...	...	...	...	Rs 2-00 to 5-00
ISI Handbook, 1972	...	...	...	...	Rs 20-00

## INDIAN STANDARDS INSTITUTION

Manak Bhavan, 9 Bahadur Shah Zafar Marg, NEW DELHI 110001

Telephone : 27 01 31 (20 lines)

Telegrams : Manaksanstha

### Branch Offices:

Telephone

'Sadhna', Nurmohamed Shaikh Marg, Khanpur, AHMEDABAD 380001	2 03 91
F Block, Unity Bldg, Narasimharaja Square, BANGALORE 560002	2 76 49
534 Sardar Vallabhbhai Patel Road, BOMBAY 400007	35 69 44
5 Chowringhee Approach, CALCUTTA 700013	23-08 02
Kothi No. 90, Sector 18 A, CHANDIGARH	
5-8-56/57 Nampally Station Road, HYDERABAD 500001	45 7 11
117/418 B Sarvodaya Nagar, KANPUR 208005	82 72
54 General Patters Road, MADRAS 600002	8 37 81
B.C.I. Bldg ( Third Floor ), Gandhi Maidan East, PATNA 800004	2 56 55